arXiv:1602.02116v3 [math.AC] 13 Sep 2016

A NOTE ON THE SUBADDITIVITY OF SYZYGIES

SABINE EL KHOURY AND HEMA SRINIVASAN

Abstract. Let $R = S/I$ be a graded algebra with t_i and T_i being the minimal and maximal shifts in the minimal graded free resolution of R at degree i. We prove that $t_n \leq t_1 + T_{n-1}$ for all n. As a consequence, we show that for Gorenstein algebras of codimension h, the subadditivity of maximal shifts T_n in the minimal graded free resolution holds for $n \geq h-1$, i.e we show that $T_n \leq T_a + T_{n-a}$ for $n \geq h-1$.

1. Introduction

Let $R = S/I$ be a standard graded algebra, with S a polynomial ring over a field k. Denote by $(\mathbb{F}, \partial)$ a minimal graded free resolution of R over S with $\mathbb{F}_a = \oplus_j S(-j)^{\beta_{aj}}$. For each a, denote by $T_a(\mathbb{F})$ and $t_a(\mathbb{F})$ the maximal and minimal shifts in the resolution $\mathbb{F}$. In other words,

$$t_a(\mathbb{F}) = min\,\{j : \beta_{aj} \neq 0\}$$

$$T_a(\mathbb{F}) = max\,\{j : \beta_{aj} \neq 0\}.$$

$\mathbb{F}$ is said to satisfy the *subadditivity* condition for maximal shifts if $T_{a+b}(\mathbb{F}) \leq T_a(\mathbb{F}) + T_b(\mathbb{F})$, for all a and b. If $\mathbb{F}$ is the minimal graded free resolution of R over S, then we write t_a and T_a for $t_a(\mathbb{F})$ and $T_a(\mathbb{F})$.

General bounds for T_n in terms of T_1 were established for all n, by Bayer-Mumford and Caviglia-Sbarra in [BM] and [CaS]. They prove that $T_n \leq (2T_1)^{2^{m-2}} - 1 + n$ where $m = \text{rank}_k S_1$. Subadditivity for maximal shifts in minimal graded free resolutions has been studied by many authors [ACI2], [EHU], [HS], [Mc]. Subadditivity for maximal shifts has been established for the highest non vanishing Tor, $Tor_p(R,k)$. It was shown that $T_p \leq T_1 + T_{p-1}$ for all graded algebras where $p = \text{projdim} R$ [HS, Corollary 3], and that $T_p \leq T_a + T_{p-a}$ when R is of depth zero and of dimension ≤ 1 [EHU, Corollary 4.1]. In [ACI2], Avramov, Conca and Iyengar consider the situation when $R = S/I$ is Koszul and show that $T_{a+1}(I) \leq T_a + T_1 = T_a + 2$ for $a \leq$ height (I). Further, they prove that if S is Cohen-Macaulay, $reg^R_{h+1}(k) = 0$ and $\binom{a+b}{a}$ is invertible in k, then $T_{a+b} \leq T_a + T_b + 1$. For monomial ideals, Herzog and Srinivasan show that $T_{a+1} \leq T_a + T_1$, for all a [HS, Corollary 4].

It is known that the minimal graded free resolution of graded algebras may not satisfy the subadditivity for maximal shifts as shown by the counter example in [ACI2]. However, no counter examples are known for monomial ideals. In fact, it can be easily seen that the subadditivity for maximal shifts does hold for algebras generated by stable monomial or square free strongly stable monomial ideals.

In this paper, we consider the minimal and maximal degrees of the syzygies and show that for any n we have $t_n \leq t_1 + T_{n-1}$. As a consequence, we prove for Gorenstein algebras S/I that $T_{a+b} \leq T_a + T_b$ for $a + b \geq s - 1$ where $s =$ height $I =$ projdim R. The question remains if the subadditivity for maximal shifts holds for all Gorenstein algebras. It is particularly interesting since the simplest counter example to whether the subadditivity holds, is a codimension four almost complete intersection linked to a codimension four Gorenstein algebra.

2. Syzygies of Gorenstein Algebras

Let $S = k[x_1, \ldots, x_d]$ and I a homogeneous ideal in S. Let $\mathbb{F}$ be the minimal graded free resolution of $R = S/I$

$$\mathbb{F} : 0 \to F_s \to \ldots \to F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} S$$

where $F_a = \oplus_j S(-j)^{\beta_{aj}}$, $t_a = min\,\{j : \beta_{aj} \neq 0\}$ and $T_a = max\,\{j : \beta_{aj} \neq 0\}$. We prove a general result for the syzygies of homogeneous algebras.

Theorem 2.1. *$R = S/I$ be a graded algebras with t_i and T_i being the minimal and maximal shifts in the minimal graded free S-resolution of R at degree i, then $t_n \leq t_1 + T_{n-1}$, for all n.*

Proof. We show the theorem by induction on n, where n is the n^{th} step of the resolution.
For $n = 1$, $T_0 = 0$ and hence $t_1 = t_1$. We need to prove the theorem for $1 < n \leq s$, where

$$\mathbb{F} : 0 \to F_s \to \ldots \to F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} S$$

is the graded resolution of R. Let $I = (g_1, \ldots, g_{\beta_1})$ where $\{g_1, \ldots, g_{\beta_1}\}$ is a set of minimal generators of I and $F_i = \bigoplus_{j=1}^{\beta_i} Sf_{ij}$ with $t_1 = \deg f_{11} \leq \deg f_{12} \leq \cdots \leq \deg f_{1\beta_1} = T_1$. Suppose $\partial_1(f_{1j}) = g_j$ and $\partial_n(f_{nt}) = \sum_{i=1}^{\beta_{n-1}} r_{ti} f_{(n-1)i}$ for all n.
We see that $Z(f_{11}, f_{1\beta_1}) = g_1 f_{1\beta_1} - g_{\beta_1} f_{11}$ is a non zero second syzygy. Hence, there exists an element $f_{11} * f_{1\beta 1} \in F_2$ of degree $t_1 + T_1$. This implies that $t_2 \leq T_1 + t_1$, and the theorem is true for $n = 2$.

Assume we have constructed by induction an element $f_{11} * f_{n-2i} \in F_{n-1}$ for all $1 \leq i \leq \beta_{n-2}$ such that

$$\partial_{n-1}(f_{11} * f_{(n-2)i}) = \partial_1(f_{11}) f_{(n-2)i} - f_{11} * \partial_{n-2}(f_{(n-2)i}),$$

and $t_{n-1} \leq t_1 + T_{n-2}$ for $4 \leq n \leq s$. Let

$$f_{11} * f_{(n-2)i} = \sum_{j=1}^{\beta_{n-1}} s_{ij} f_{(n-1)j}.$$

We also have $\partial_{n-1}(f_{(n-1)t}) = \sum_{i=1}^{\beta_{n-2}} r_{ti} f_{(n-2)i}$ for all $1 \leq t \leq \beta_{n-1}$. Consider

$$Z(f_{11}, f_{n-1t}) = \partial_1(f_{11}) f_{n-1t} - \sum_{i=1}^{\beta_{n-2}} \sum_{j=1}^{\beta_{n-1}} r_{ti} s_{ij} f_{(n-1)j}.$$

$Z(f_{11}, f_{n-1t})$ is a cycle in F_{n-1}, since

$$\begin{aligned}
\partial_{n-1}(Z(f_{11}, f_{(n-1)t})) =& \partial_1(f_{11})\partial_{n-1}(f_{(n-1)t}) - \sum_{i=1}^{\beta_{n-2}} r_{ti}\partial_{n-1}(\sum_{j=1}^{\beta_{n-1}} s_{ij}f_{(n-1)j}) \\
=& \partial_1(f_{11}) \sum_{i=1}^{\beta_{n-2}} r_{ti}f_{(n-2)i} - \sum_{i=1}^{\beta_{n-2}} r_{ti}\partial_{n-1}(f_{11} * f_{(n-2)i}) \\
=& \sum_{i=1}^{\beta_{n-2}} \left[\partial_1(f_{11})r_{ti}f_{(n-2)i} - \partial_1(f_{11})r_{ti}f_{(n-2)i} + r_{ti}f_{11} * \partial_{n-2}(f_{(n-2)i})\right] \\
=& \sum_{i=1}^{\beta_{n-2}} r_{ti}f_{11} * \partial_{n-2}(f_{(n-2)i}) \\
=& f_{11} * \partial_{n-2}(\sum_{i=1}^{\beta_{n-2}} r_{ti}f_{(n-2)i}) \\
=& f_{11} * \big(\partial_{n-2} \circ \partial_{n-1}(f_{(n-1)t})\big) \\
=& f_{11} * 0 \\
=& 0.
\end{aligned}$$

Thus, for every t, $1 \leq t \leq \beta_{n-1}$, there exists an element $f_{11} * f_{(n-1)t} \in F_n$ of the same degree as $\deg Z(f_{11}, f_{(n-1)t})$ that is mapped by ∂_n onto $Z(f_{11}, f_{(n-1)t})$. This means there is an element $f_{11} * f_{(n-1)t} \in F_n$ of degree $t_1 + deg f_{(n-1)t} \leq t_1 + T_{n-1}$ such that $\partial_n(f_{11} * f_{(n-1)t}) = Z(f_{11}, f_{(n-1)t})$. However, the cycles $Z(f_{11}, f_{n-1t})$ can be zero for all t! For that, we show that there is at least one t such that one of the cycles $Z(f_{11}, f_{n-1t})$ is not identically zero.
Suppose that $Z(f_{11}, f_{n-1t}) = 0$ for all t, then $\partial_1(f_{11})f_{n-1t} = \sum_{i=1}^{\beta_{n-2}} \sum_{j=1}^{\beta_{n-1}} r_{ti}s_{ij}f_{n-1j}$. So for all t, we have

$$\sum_{i=1}^{\beta_{n-2}} \sum_{j=1}^{\beta_{n-1}} r_{ti}s_{ij} = \begin{cases} 0 & j \neq t \\ \partial_1(f_{11}) & j = t. \end{cases}$$

By setting $\bar{r} = \bar{r}_{n-1} = (r_{ti})_{\beta_{n-1} \times \beta_{n-2}}$ and $\bar{s} = (s_{ij})_{\beta_{n-2} \times \beta_{n-1}}$, we get $\bar{r}\bar{s} = \partial_1(f_{11})I$ and hence the rank $\bar{r}_{n-1}\bar{s} = \beta_{n-1}$. By the exactness of the resolution, we have rank $\bar{r}_n +$ rank $\bar{r}_{n-1} = \beta_{n-1}$ where $\bar{r}_n$ is the matrix representing ∂_n. Since $\partial_n \neq 0$, this implies that rank $\bar{r}_{n-1} < \beta_{n-1}$ which is a contradiction.

Hence, there exists a non zero cycle in F_{n-1}, and we get $t_n \leq t_1 + T_{n-1}$. ■

The following remark is an easy consequence of the duality of the minimal graded free resolution for Gorenstein algebras.

Remark 2.2. If $R = S/I$ is a Gorenstein algebra, with height $I = h$, then $T_h \leq T_a + T_{h-a}$.
Since $c = T_h = t_h$, then by the duality of the minimal graded free resolution $\mathbb{F}$ we get $c - t_{h-a} = T_a$ for all $a = 1, \cdots h-1$. This implies that $c = T_a + t_{h-a} \leq T_a + T_{h-a}$.

Theorem 2.3. *For any graded Gorenstein algebra S/I of codimension h, we have $T_{h-1} \leq T_a + T_{h-1-a}$. Thus, $T_n \leq T_a + T_{n-a}$ for $n \geq h-1$.*

Proof. Since S/I is Gorenstein, then $T_{h-1} = T_h - t_1$ and $T_{h-1-a} = T_h - t_{a+1}$. So, $T_{h-a-1} = T_h - t_{a+1} \geq T_h - (t_1 + T_a)$ by theorem 2.1. So, $T_{h-a-1} \geq T_h - t_1 - T_a = T_{h-1} - T_a$ and hence $T_{h-1} \leq T_a + T_{h-a-1}$ as desired. ∎

Definition 2.4. A minimal graded free resolution is said to be pure if, for all a, F_a is generated in one degree. Hence the minimal graded free resolution is pure if it has the following shape

$$\mathbb{F} : 0 \to S(-j_s)^{\beta_s} \to \ldots \to S(-j_2)^{\beta_2} \xrightarrow{\partial_2} S(-j_1)^{\beta_1} \xrightarrow{\partial_1} S$$

Corollary 2.5. *For any graded algebra S/I with a pure resolution, we get $T_n \leq T_1 + T_{n-1}$.*

Proof. When the resolution is pure we have $t_a = T_a$ for all a, and hence by theorem 2.1 we get the result. ∎

Remark 2.6. As a consequence, if S/I is a Gorenstein pure algebra of codimension 4 or 5, then its minimal graded free resolution satisfies the subadditivity condition.

Remark 2.7. As the following three examples show, the inequality can be tight even for codimension four Gorenstein algebras with pure resolutions.

Example 2.8. Let $S = k[x, y, z, w]$ and $I = (w^2, z^2, y^2 - wz, wx^2, x^2z, x^2y + wyz, x^3 + wxz + wyz)$ a height four Gorenstein ideal. We have $T_4 = 8$, $T_3 = 6$ $T_2 = 4$ and $T_1 = 3$. We get $T_4 = 2T_2 \leq T_3 + T_1$ and $T_3 \leq T_2 + T_1$. ∎

Example 2.9. Let $S = k[x, y, z, w]$ and $I = (w^3, w^2z, w^2y, wz^2, wyz - w^2x, wy^2, z^3 + wxy, yz^2, xz^2, y^2z, xyz - wx^2, x^2z, y^3 - 2wxz, xy^2, x^2y + w^2x, x^3 + wxz - w^2x)$ a height four Gorenstein ideal generated in degree three and having a pure resolution. Then the minimal graded free resolution of S/I satisfies the subadditivity condition. We have $T_4 = 8$, $T_3 = 5$, $T_2 = 4$ and $T_1 = 3$. Then we get $T_4 = 2T_2 = T_3 + T_1$, $T_3 \leq T_2 + T_1$ and $T_2 \leq 2T_1$. ∎

Example 2.10. Let $S = k[x, y, z, w, r, s, t]$ and $I = (rt, tz, ty - st, tx, sw - s^2, sz, sy, sx, rw - st - t^2, rz, yr + tw, rx - s^2, w^2 + st, wz, wy - s^2, wx, yz - z^2 + r^2 + rs + tw - st - t^2, xz - rs - tw + st + t^2, y^2, xy - z^2 - r^2 + rs + tw - st - t^2, x^2 + rs + tw - st - t^2)$ a height seven Gorenstein ideal with $T_7 = 10$, $T_6 = 8$, $T_5 = 7$, $T_4 = 6$, $T_3 = 5$, $T_2 = 3$ and $T_1 = 2$. Then we get $T_7 = T_6 + T_1 = T_5 + T_2 \leq T_4 + T_3$ and $T_6 \leq T_5 + T_1 = T_4 + T_2 \leq 2T_3$.

∎

References

[ACI1] L.L. Avramov, A. Conca, and S. B. Iyengar, *Free resolutions over commutative Koszul algebras*, Math. Res. Lett. **17** (2010), no. 2, 197–210.

[ACI2] L.L. Avramov, A. Conca, and S. B. Iyengar, *Subadditivity of Syzygies of Koszul algebras*, Math. Ann. **361** (2015), no. 1-2, 511–534. 1

[BM] D. Bayer, D. Mumford, *What can be computed in algebraic geometry?, in Computational Algebraic Geometry and Commutative Algebra (Cortona, 1991)*, Sympos. Math., vol. XXXIV (Cambridge University Press, Cambridge, 1993), pp. 1–48 1

[EHU] D. Eisenbud, C. Huneke and B. Ulrich, *The regularity of Tor and graded Betti numbers*, Amer. J. Math. **128** (2006), no. 3, 573–605. 1

[FG] O. Fernändez-Ramos, P. Gimenez, *Regularity 3 in edge ideals associated to bipartite graphs*, J. Algebraic Combin. 39 (2014), no. 4, 919–937.

[CaS] G. Caviglia, E. Sbarra, *Characteristic-free bounds for the Castelnuovo-Mumford regularity*, Compos. Math. **1**41(6) (2005), 1365–1373 . 1

[HS] J. Herzog, H. Srinivasan, *A note on the subadditivity problem for maximal shifts in free resolutions*, Preprint 2013, arXiv:1303.6214. 1

[Mc] J. McCullough, *A polynomial bound on the regularity of an ideal in terms of half the syzygies*, Math. Res. Lett. **19** (2012), no. 3, 555–565 1

Department of Mathematics, American University of Beirut, Beirut, Lebanon.
E-mail address: se24@aub.edu.lb

Department of Mathematics, University of Missouri, Columbia, Missouri-65211, USA.
E-mail address: srinivasanh@missouri.edu